



ORIGINAL CONTRIBUTION


The Relationship of Perceived Stress, Emotional Intelligence and Resilience with Subjective Well Being among Nurses

Mahekamil Latif Abbasi ¹, Zarsha Khan ², Tehreem Islam ³, Nasreen Rafique Kausar ⁴, Museera Aymen ^{5*}
1,2,3,4,5 Shifa Tameer-e-Millat University, Islamabad, Pakistan

Abstract— The study investigates the association between nurses' subjective well-being, emotional intelligence, resilience, and perceived stress. The purpose of the study is to better understand how nurses' subjective well-being effected by stress, emotional intelligence and resilience. The study conducted in Islamabad and Rawalpindi. The data was collected from public and private hospital of Islamabad. Study started in November 2022 and end in June 2023. Correlational research design were used A broad collection of nurses from various healthcare settings were gathered with a sample size $N = 300$ (calculated by Rao soft), with ages ranging from (20-50 years). The background details of nurses were collected with the use of demographic form. The Perceived Stress Scale (PSS), Schutte Self-Report Emotional Intelligence Test, Brief Resilience Scale (BRS) and Satisfaction with Life Scale (SWLS) were used. The results revealed a significant relationship between perceived stress, emotional intelligence, and job experience with the subjective well-being of nurses, predicting the role of perceived stress, emotional intelligence, resilience, and job experience on the well-being of nurses. The values ($\beta = .41$, $p = 0.00$) showed a 41% variance ($R^2 = 0.41$). The research on the connections between perceived stress, emotional intelligence, resilience, and subjective well-being among nurses, in particular, offers important new understandings of the variables affecting the general well-being of this crucial healthcare profession.

Index Terms— Perceived stress, Emotional intelligence, Resilience, Subjective well-being, Nurses

Received: 03 February 2023; **Accepted:** 31 April 2023; **Published:** 28 July 2023



Introduction

In the intricate tapestry of healthcare, nurses stand out as the caring threads that weave patients, families, and healthcare professionals together in the complex fabric of healthcare. Beyond their clinical expertise, nurses exhibit the best of humanity by providing comfort, assistance, and healing to those in need (Choo et al., 2010). With their unshakable commitment, compassionate care, and relentless advocacy, nurses play a crucial part in the healthcare industry, influencing patients' lives and fostering general well-being.

The nursing profession has grown into an indispensable pillar of contemporary healthcare, with roots in centuries of tradition and innovation. By acting as the link between medical expertise and the human condition, nurses constitute the backbone of patient-centered care. Their presence offers security, stability, and a caring touch that bridges the gap between disease and healing, from the busy emergency rooms to the calm hallways of long-term care institutions (Raatikainen, 1997).

Additionally, nurses easily work in interdisciplinary teams with doctors, therapists, chemists, and other healthcare specialists. They serve as the central figure in care coordination, coordinating and integrating many areas of care to enable a comprehensive strategy that

*Email: museeraaymen@gmail.com

improves patient outcomes. Their proficiency in care management, medication administration, and patient monitoring guarantees the efficient functioning of healthcare facilities, fostering an atmosphere that is supportive of healing and rehabilitation.

In the healthcare industry, nurses are the pinnacle of compassion, skill, and unshakable commitment. They represent the very best of humanity in healthcare, playing a role that goes far beyond mere medical knowledge. Through their all-encompassing care, advocacy, emotional support, and teamwork, nurses light the way to recovery and help patients and their families through the maze of disease (Al-Dalaeen et al., 2023; Bonsall & Cheater, 2008). However, their altruistic dedication frequently has a cost, though, as the constant stresses and demands of the job can result in burnout, which is a state of physical, emotional, and mental tiredness. Even the most resilient people can be tested to the breaking point in the stressful profession of nursing, despite it being intrinsically honorable and fulfilling (Cheater, 2008).

The well-being of nurses is negatively impacted by the long hours, large patient loads, difficult cases, and the constantly shifting healthcare environment. Burnout among nurses has effects that go beyond the person. Patient care quality may worsen when nurses become less engaged and satisfied with their work. Burnout can impair one's ability to think critically, make decisions, and pay attention to detail, which can result in mistakes in patient care and medical blunders. Burnout can affect the nurse-patient relationship, obstructing good communication and delaying the healing process. Burnout also harms nurses' personal lives, which exacerbates the cycle of stress and burnout by causing strained relationships, poorer work performance, and more absences (Spence Laschinger & Fida, 2014).

Literature Review

Perceived stress is the cognitively processed emotional response to the target event rather than the possibly continuous life event that occurs to the individual (Cohen et al., 1983). Emotions are a part of our daily life and managing those emotions whether negative or positive is important for us to maintain our everyday functioning and perform better in our social roles (Scheibe et al., 2018).

Factors associated with job-related stress among nurses have frequent exposure to adverse events at work, increase in workload, shortage of nurses at hospitals, emotional demands of patients and their families, compassion fatigue, and compassion satisfaction (Ekedahl & Wengström, 2007, Stamm et al., 2009).

The nature of a nurse's work can sometimes be stressful. Nurses frequently have little autonomy and control over their jobs, are transferred between patient care units, have poor communication among members of the health care team, and frequently deal with demanding or recalcitrant family members of patients. Work-related stressors like these heighten nurses' feelings and perceptions of being overworked and stressed (Jordan et al., 2016).

Emotional intelligence is the ability to identify, understand and balance emotions and then utilize them in one's life (Morris & Wakefield, 2008). The human brain interprets emotional information from the environment diversely at an individual level. The exposition of grasping the idea of emotional intelligence varies from individual to individual, hence called emotional intelligence (Mayer, 1997); the power to access and produce feelings when they help cognition; the ability to understand affect-laden knowledge utilize it; and the power to control the emotions to aid welfare (Mayer & Salovey 1997).

The four-branch model of emotional intelligence outlines four areas of abilities or skills that jointly represent many different aspects of emotional intelligence. The abilities to effectively identify one's own and others' emotions, use emotions to help thinking, comprehend emotional meanings, and control emotions are explicitly referred to in this concept as emotional intelligence (Ali et al., 2010; Salovey & Mayer, 1997). It has a significant impact on people's job performance and also on one's emotional capacity which plays an important role in health care workers like nurses treating their patients and bearing the emotional burden (Morris & Wakefield, 2008). Emotional intelligence is one of the most important aspects that help improve psychological health (Cooper & Sawaf, 1997).

Studies have shown that emotional intelligence is not just a preventive element against negative sentiments, but is also connected to psychological health (Mayer et al. 1999, Brackett & Mayer 2003; Brackett et al. 2006; Brackett et al. 2011; Zeidner et al. 2012). In this light emotional intelligence has proven to be connected to many essential real-life upshots such as positive mental health and subjective well-being (Mayer et al. 2008). Researchers indicate that elevated emotional intelligence is recorded by ability tests and is linked to well-being indicators (Brackett & Mayer 2003, Extreme et al. 2011).

Positive psychology has attracted scientists over the last ten years (Seligman & Csikszentmihalyi 2000, Seligman 2003). Theoretical models of positive psychology could be enhanced by the incorporation of dissimilarities in the processing of affective information because the sharp and clever use of emotions is thought to be vital for one's psychological adaptation and welfare (Brackett & Mayer 2003).

A recent study on female nursing students showed higher emotional intelligence levels which are associated with less perceived stress and higher levels of life satisfaction and happiness than the general population (Aranda, Extreme, & Galán, 2014). Nurses with high emotional intelligence have less emotional weariness and psychosomatic symptoms; they have better emotional health; they derive more satisfaction from their actions (both at work and home); and they have stronger relationships with colleagues (Towell et al, 2013). Nurses with strong emotional intelligence have been found to have good health and well-being (Hao Hung et al, 2022).

Resilience is the quality of an individual to adapt to difficult situations and come out of them successfully. Resilience is obtained when one depicts emotional and behavioral flexibility to adjust to difficult life experiences regarding internal and external demands (Norman Garmezy, 1991). Research has shown that an individual with a greater sense of emotional intelligence shows the traits of resilience and also greater self-esteem (Masten et al, 2001). Resilience plays an important role in the lives of nurses as they come by with numerous difficulties and adverse cases of their patients (Trigueros et al., 2020). Resilience comes with the existence of three main stages.

Nurses are frontline workers and they are providing quality care to the patient, establishing an effective rapport, and empathizing with their patients while managing their own emotions (Evans & Allen, 2002). Resilience is one's successful adaptation to any stressful situation or adverse events in life (Masten et al., 1994). More resilient individuals can easily recover from negative events or remain healthy after adverse events in their life as compared to people who are more vulnerable to negative events (Hurley et al., 2004). The concept of subjective well-being given by Diener (2009), emphasizes the importance of happiness and well-being for the mental health of the individual. Individuals' subjective well-being is associated with the cognitive evaluation of one's own life as well as the experience of positive and negative emotions and events of one's own life (Diener et al., 2008).

Subjective well-being is the self-reported evaluation of one's quality of life, using one's happiness as the criterion which is a general key indication of individual quality of life (Lucas 2002); Preventing, alleviating, and lowering burnout all have a positive impact on people's subjective well-being (Diener & Lucas 2002). The circumstance theory of subjective well-being states that the well-being of an individual is the result of different circumstances in one's life and positive and negative events in everyday life (Diener et al, 2009). According to dispositional views, subjective well-being is essentially the product of biological or temperamental variables that impact how we interpret and judge life circumstances and events, rather than actual conditions themselves (James E Maddux, 2020). Optimism, life satisfaction, and good feelings are all examples of subjective well-being (Diener et al., 2009).

Subjective well-being has been shown to have an impact on many aspects of life, including work performance and mental health (Diener & Ryan, 2009). Having a high level of subjective well-being is desirable in people, achieving a happy life and ways to increase overall well-being seems more important in people now (Milosevic et al., 2011). Subjective well-being is not only beneficial to one's mental and physical health but also contributes to health organizations as individuals whose subjective well-being is high are more creative, stay motivated at work, and form healthy relationships in life (Gavin et al., 2014).

There is a significant positive correlation seen between subjective well-being and job satisfaction in nurses (Laring Yu et al., 2015). Nurses specifically working in ICU are more prone to experience job stress (Guo et al., 2013). Subject to high intensity of stress at work, fewer opportunities to contact the outside world, care for critically ill patients, gaps between technological advancements and nursing skills, and the incidence of death leads to stress (ChingLing Sun et al., 2013).

Due to the demanding nature of nursing, nurses frequently experience significant levels of stress, emotional weariness, and burnout. Promoting and ensuring their subjective well-being is therefore critical to their work and personal fulfillment (Diener, 2018). Zhang (2020) studied that those nurses who reported higher levels of subjective well-being showed better levels of job satisfaction, lower levels of stress, and higher levels of involvement in their work. These favorable outcomes help not just the nurses themselves, but also patient care and organizational performance (Zhang et al., 2020).

Rationale

Healthcare professionals that work in the challenging field of nursing must meet high physical, mental, and emotional demands. In addition to being important for their satisfaction, nurses' well-being also plays a key role in delivering high-quality patient care (Brennan, 2017). A key determinant of the nurses' overall quality of life is subjective well-being, which includes an individual's cognitive and affective assessments of their general life satisfaction and enjoyment. Investigating the variables that affect nurses' subjective well-being is crucial. The purpose of this study is to examine, from a human-centric standpoint, the relationship between perceived stress, emotional intelligence, resilience, and subjective well-being among nurses (Gurková et al., 2017).

It is crucial to comprehend the complex interactions between perceived stress, emotional intelligence, resilience, and subjective well-being in nurses. This research aims to recognize the complexity of nurses. Understanding these characteristics better would enable healthcare organizations, policymakers, and educators to collaborate and develop environments that support nurses' well-being, thereby benefiting both medical staff members and the patients they look after. This research aims to acknowledge the significance of nurses' subjective experiences, emotions, and well-being (Román et al., 2017). The most detrimental type of stress to a person's mental and physical health is psychological stress. Reduced contentment is associated with intense stress (Suleman, 2018). It recognizes that internal processes such as stress assessment, emotional intelligence, and resilience, as well as external elements like workload and organizational support, also have an impact on nurses' well-being. We aim to provide a thorough knowledge of the complex interactions that affect nurses' subjective well-being by looking at these internal elements.

Research Objectives

1. To explore the relationship between perceived stress and subjective well-being of nurses working in public and private hospitals of Islamabad/Rawalpindi.
2. To find out the relationship between emotional intelligence and the subjective well-being of nurses
3. To explore the role of resilience as a correlate of nurses' subjective well-being.
4. To find out the effect of demographic variables such as gender, marital status, residence, area, and income on nurses' well-being.

Methods

Research Design

A quantitative research method using multiple linear regression analysis was used to assess the relationship between perceived stress, emotional intelligence, and resilience with subjective well-being among nurses of age 20-50 years. The data was collected purposely using quantitative measurement tools.

Sample

The g-power sample size calculator was used to calculate the sample size for this study. The total number of participants was 300. Data was collected using a purposive sampling technique from twin cities Rawalpindi and Islamabad. The age range of the respondents was 20-50 years. The data was collected manually through printed forms.

Sample Selection Criteria

Inclusion: The inclusion criteria for participant selection were adult nurses, ages ranging from 20-50 years. This included both genders (males and females) residing within Pakistan. Participants had an equal chance to be selected because of purposive sampling.

Exclusion: Nurses less than the age of 20 were not included in the research.

Demographic Sheet

The demographic sheet contained the relative information about the participant's age, gender, job experience in years, marital status, and number of family members.

Perceived Stress Scale

The perceived stress scale is a 10-item questionnaire developed by Cohen et al.1983. It is used to measure perceived helplessness and lack of self-efficacy in an individual which results in a stressful state in young people and adults aged 12 and above, Cohen reported that the reliability of this scale is 0.78. According to our results Cronbach alpha based on standardized items is 0.69.

Schutte Self Report Emotional Intelligence Test (SSEIT)

Schutte self-report emotional intelligence test (SSEIT) a 33 item scale is given by Salovey and Mayer, 1990. The scale uses four subscales; emotion perception, utilizing emotions, managing self-relevant emotions, and managing others' emotions. The reliability rating of this scale is 0.90 for emotional intelligence given by Mayer.

Brief Resilience Scale

Brief Resilience Scale was developed by Smith, 2008 to assess an individual's perceived ability to recover or bounce back from stress. The scale was designed to assess a unitary construct of momentary resilience, using items that were both positively and negatively worded. Brief Resilience Scale has a reliability of 0.71 given by Smit (2008).

The Satisfaction with Life Scale

The satisfaction with life scale is a five item questionnaire developed by Diener, 1985 which measures the global cognitive judgment of one's life satisfaction. The reliability of this scale is 0.82 according to Diener.

Procedure

The topic of this study was approved by the Departmental Research advisory committee of Shifa Tameer-e-Millat University, after which a non-objection certificate from the university was given concerning reaching out to the authorities of the hospitals who were approached for the collection of data, and then was debriefed about the study. After the collection of data was permitted, data was gathered from the hospitals of Rawalpindi and Islamabad. The participants (nurses) were also debriefed about the purpose of the study. Nurses who wished to take part were separated from the ones who did not wish to be a part of this study. The participants who agreed to participate were then given an informed consent form. After this, they were given five questionnaires were used in survey.

Results

The present study investigated the relationship between perceived stress, emotional intelligence, and resilience with Subjective well-being among nurses. The analyses were done through the SPSS version 25. Perceived Stress Scale, Schutte Self-Report Emotional Intelligence Test (SSEIT), Brief resilience scale, and satisfaction with Life Scale SWLS were used to measure Perceived stress, emotional intelligence, and resilience with subjective well-being among nurses.

Table I
Descriptive Characteristics for Socio-Demographics Variable (N = 300)

	Variables	f	%
Gender	Male	104	35
	Female	193	65
Job experience in years	1-10	173	57
	10-20	4	1
	20-30	123	41
Age	20-30	190	63
	30-40	106	35
	40-50	4	1
Marital Status	Single	159	53
	Married	141	47
No of family member	1-5	174	58
	5-10	119	39
	10-15	7	3

Note: f = frequency, % = percentage

The table above shows the frequencies of all socio-demographic variables of the participant $n = 300$. 104(34.7%) were males and 193 (%64.3) were females. In the data set three categories of age were found; $f = 190(63\%)$ participants fall in the range of 20-30, $f = 106(35\%)$ lie in the range of 30-40 and remaining $f = 4(1\%)$ lie in the 40-50 category. 159(53%) were single whereas 141(47%) were married. 173(57%) have 1-10 years of job experience, 4 participants have 10-20 years of job experience and 123(41%) have 20-30 years of job experience.

Table II
Correlation Matrix of all Scales (N = 300)

S No	Variables	n	M	S.D	1	2	3	4
1	Perceived stress	300	30	4.2	-	-	-	-
2	Emotional intelligence	300	116	16.6	-.110**	-	-	-
3	Resilience	300	18.1	3.5	-.371	.081	-	-
4	Subjective well being	300	17.1	5.6	.341**	-.053	-.174**	-

Note: M = Mean, n = Total Participant, SD = Standard Deviation

Correlation analyses were used to analyze the relationship between perceived stress, emotional intelligence, resilience, and subjective well-being among nurses. The result showed that emotional intelligence and perceived stress have a highly statistically significant positive correlation $p < 0.01$. It concluded that with the increase in perceived stress, there is a significant increase in emotional intelligence. Resilience is non-significant.

Table III

Multiple Linear Regressions Predicting Wellbeing from Perceived Stress, Emotional Intelligence, Resilience, And Job Experience ($N = 300$)

Predictors	SE	B	p	ΔF	R
				15.01	.41
Constant	4.0	6.7	0.00		
1	Perceived stress,	.077	.426	0.00	
2	Emotional intelligence	.020	-.087	0.00	
3	Resilience	.091	.047	.97	

Note. B = unstandardized beta, S.E = standard error, p = significance level

Multiple Linear regression was used to assess the predicting role of perceived stress, emotional intelligence, resilience, and job experience on the well-being of nurses. The table shows predicting the role of perceived stress, emotional intelligence, resilience, and job experience on the well-being of nurses. The values ($\beta = .41$, $p = 0.00$) showed a 41% variance ($R^2 = 0.41$) so perceived stress, job experience, and emotional intelligence predict subjective well-being in nurses. For every one-unit increase in perceived stress, there is a 41-unit increase in subjective well-being. There was no other significant variance found.

Discussion

The purpose of the research study is to explore the relationship between perceived stress, Emotional Intelligence, and Resilience with subjective well-being among nurses working in public and private hospitals of Islamabad/Rawalpindi. The sample population encompassed nurses with ages ranging from 20-50 years. Participants were recruited using purposive sampling techniques from the cities of Rawalpindi and Islamabad.

The emotional intelligence predicts positive subjective well-being in nurses. According to our analysis, our hypothesis is true and is concurrent with the research. To increase nurses' positive subjective well-being, emotional intelligence is crucial (Goleman, 2006). Self-awareness, or the capacity to recognize and comprehend one's own emotions, strengths, and limitations, is a component of emotional intelligence. Nurses who are very self-aware are better able to control their emotions and deal with difficult situations. They can recognize their stress triggers, admit when they require assistance, and practice self-care (Codier et al., 2012). This self-awareness enables nurses to manage their emotional state proactively, maintaining positive subjective well-being. Effective emotion regulation is made possible for nurses by emotional intelligence. It requires being able to identify and control one's own and other people's emotions. Nursing professionals can react to pressures, disappointments, and challenging situations more effectively by comprehending and controlling their emotions. To keep a positive emotional state, they can use techniques like deep breathing, mindfulness, and reframing unfavorable ideas (Atanes AC; Andreoni S; Hirayama MS; Montero-Marin J; Barros VV; Ronzani TM; Kozasa EH; Soler J; Cebolla A; Garcia-Campayo J; Demarzo MM). Effective emotion control helps people feel better and experience less stress. Emotional intelligence enables nurses to effectively control their emotions. It necessitates being able to recognize and manage one's own emotions as well as those of others. Understanding and managing their emotions helps nurses respond to demands, disappointments, and difficult situations more skillfully. They can practice methods like deep breathing, mindfulness, and reframing adverse thoughts to maintain a positive emotional state. People who can effectively control their emotions feel better and are under less stress.

The resilience correlates with the subjective well-being of nurses. Resilience is a crucial element that has a favorable correlation with nurses' subjective well-being (Al-Turki et al., 2010). To effectively manage stress, adapt to changes, control emotions, solve problems, maintain work-life balance, and feel a feeling of mastery and growth, nurses must develop resilience. Together, these elements support their health and ability to offer their patients high-quality care.

According to a study by Yuan Qin (2023), an individual's emotional intelligence has an impact on their level of life satisfaction. According to the positive psychology perspective, nursing managers should provide nurses with a nurturing work atmosphere to boost their confidence and resilience, which will ultimately raise their degree of life satisfaction. This study's findings corroborate our own, and it is amply demonstrated that nurses who report feeling better about their subjective well-being have higher levels of life satisfaction. Zakeri et al. (2021) found that nurses who were dissatisfied with their lives were up to 2.4 times more likely to suffer from psychological diseases. According to John D. Mayer, Salovey, and Caruso (2004), strong emotional intelligence is associated with a greater propensity to be drawn to socially involved professions including nursing, medicine, psychology, and teaching. According to Cooper & Sawaf (1997), one of the key factors in enhancing psychological wellness is emotional intelligence. According to Mayer et al. (1999), emotional intelligence is not just a factor in preventing the development of unpleasant feelings (Adriaenssens et al., 2015).

Limitations

The study is only focused on working nurses who are between the ages of 20 and 50. Because of the short sample size, it was unable to distinguish between gender and city differences. The data was only gathered from Islamabad and Rawalpindi, two cities that are both in Punjab and one of which is the federal capital of Pakistan. There were no omitted cities. Measures that nurses self-report were subject to response biases and may have not correctly reflected their actual experiences or behaviors.

Future Discussion

On nursing students, additional research can be done. In Pakistan, nurses and nursing students can both be researched in relation to other variables. Future studies might include an equal number of men and women. The sample size can also be expanded to include Pakistan's other major cities. We can investigate the interactions between perceived stress, emotional quotient, resilience, and subjective well-being. Contextual influences include work environment, social support, and organizational culture. Having a thorough understanding of these contextual elements can help us develop supportive work settings that encourage nurses' wellbeing. We can Conduct cross-cultural research to learn how cultural differences affect how nurses experience stress in connection to their emotional intelligence, resilience, and subjective well-being. This can guide the creation of therapies that are specifically suited to a certain cultural setting by identifying culturally distinctive characteristics that affect subjective well-being.

Conclusion

The research on the connections between perceived stress, emotional intelligence, resilience, and subjective well-being among nurses, in particular, offers important new understandings of the variables affecting the general well-being of this crucial healthcare profession. The results of much research demonstrate the complex interactions between these factors and how they affect nurses' subjective well-being. Healthcare organizations should try to develop supportive work environments that encourage emotional intelligence, resilience, and subjective well-being among nurses in light of the significance of these traits. Nurses can have a better and more satisfying work experience by implementing stress management programs, giving resources for enhancing emotional intelligence abilities, and supporting resilience-building efforts. Given the importance of these qualities, healthcare organizations should seek to provide supportive work environments that foster nurses' emotional intelligence, resilience, and subjective well-being. By adopting stress management courses, providing tools for increasing emotional intelligence skills, and assisting resilience-building initiatives, nurses can have a better and more fulfilling work experience.

REFERENCES

- Adriaenssens, J., De Gucht, V., & Maes, S. (2015). Determinants and prevalence of burnout in emergency nurses: A systematic review of 25 years of research. *International Journal of Nursing Studies*, 52(2), 649–661. <https://doi.org/10.1016/j.ijnurstu.2014.11.004>
- Ali, A., Ahmad-Ur-Rehman, M., Haq, I. U., Jam, F. A., Ghafoor, M. B., & Azeem, M. U. (2010). Perceived organizational support and psychological empowerment. *European Journal of Social Sciences*, 17(2), 186-192.
- Altun, I., Goksel, E., & Kitapci, H. (2014). Subjective well-being of nurses. *Journal of the Pakistan Medical Association*, 64(12), 1434.
- Al-Dalaeen, A. S., Shoqirat, M. D. A., Alshawawreh, S., & Alzaben, M. B. L. (2023). Challenge and Hindrance Stressors and Mental Health Influencing Psychological Well Being: Moderation of Psychological Capital. *Pakistan Journal of Life & Social Sciences*, 21(1).
- Al-Turki, H., Al-Turki, R., Al-Dardas, H., Al-Gazal, M., Al-Maghrabi, G., Al-Enizi, N., & Ghareeb, B. (2010). Burnout syndrome among multi-national nurses working in Saudi Arabia. *Annals of African Medicine*, 9(4), 226. <https://doi.org/10.4103/1596-3519.70960>
- American Psychological Association. (n.d.). *Apa PsycNet*. American Psychological Association. <https://psycnet.apa.org/record/2004-10747-009>
- Atanes, A. C., Andreoni, S., Hirayama, M. S., Montero-Marin, J., Barros, V. V., Ronzani, T. M., Kozasa, E. H., Soler, J., Cebolla, A., Garcia-Campayo, J., & Demarzo, M. M. (n.d.). *Mindfulness, perceived stress, and subjective well-being: A correlational study in Primary Care Health Professionals*. BMC Complementary and Alternative Medicine.
- Bonsall, K., & Cheater, F. M. (2008). What is the impact of advanced primary care nursing roles on patients, nurses and their colleagues? A literature review. *International Journal of Nursing Studies*, 45(7), 1090–1102. <https://doi.org/10.1016/j.ijnurstu.2007.07.013>
- Brackett, M. A., Mayer, J. D., & Warner, R. M. (2004). Emotional intelligence and its relation to everyday behavior. *Personality and Individual Differences*, 36(6), 1387–1402. [https://doi.org/10.1016/s0191-8869\(03\)00236-8](https://doi.org/10.1016/s0191-8869(03)00236-8)
- Brennan, E. J. (2017). Towards resilience and wellbeing in nurses. *British Journal of Nursing*, 26(1), 43–47. <https://doi.org/10.12968/bjon.2017.26.1.43>
- Codier, E., Freitas, B., & Muneno, L. (2012). Developing emotional intelligence ability in oncology nurses: A clinical rounds approach. *Oncology Nursing Forum*, 40(1), 22–29. <https://doi.org/10.1188/13.onf.22-29>
- Diener, E., Oishi, S., & Tay, L. (2018). Advances in subjective well-being research. *Nature Human Behaviour*, 2(4), 253-260.
- Diffley, D. M., & Duddle, M. (2022). Fostering Resilience in Nursing Students in the Academic Setting: A Systematic Review. *Journal of Nursing Education*, 61(5), 229-235.
- University of West Florida. (2020) *Emotional intelligence in nursing*. Retrieved from <https://shorturl.at/klmIR>
- Görgens-Ekermans, G., & Brand, T. (2012). Emotional intelligence as a moderator in the stress-burnout relationship: A questionnaire study on Nurses. *Journal of Clinical Nursing*, 21(15–16), 2275–2285. <https://doi.org/10.1111/j.1365-2702.2012.04171.x>
- Gurková, E., Čáp, J., Žiaková, K., & Ďurišková, M. (2011). Job satisfaction and emotional subjective well-being among Slovak nurses. *International Nursing Review*, 59(1), 94–100. <https://doi.org/10.1111/j.1466-7657.2011.00922.x>
- Hakanen, J. J., Bakker, A. B., & Schaufeli, W. B. (2019). How work engagement and burnout are related to subjective health and sickness absence: A two-wave study. *Journal of Psychological Health*, 34(6), 667-686.
- Huang, H., Liu, L., Yang, S., Cui, X., Zhang, J., & Wu, H. (2019). Effects of job conditions, occupational stress, and emotional intelligence on chronic fatigue among Chinese nurses: A cross-sectional study. *Psychology Research and Behavior Management*, 351-360.
- Jefferies, D., Johnson, M., & Ravens, J. (2011). Nurturing and nourishing: The nurses' role in nutritional care. *Journal of Clinical Nursing*, 20(3–4), 317–330. <https://doi.org/10.1111/j.1365-2702.2010.03502.x>
- Jordan, T. R., Khubchandani, J., & Wiblehauser, M. (2016). The impact of perceived stress and coping adequacy on the health of nurses: A pilot investigation. *Nursing Research and Practice*, 2016.
- Mayer, J. D., Salovey, P., & Caruso, D. R. (2004). Target articles: "Emotional intelligence: Theory, findings, and implications." *Psychological Inquiry*, 15(3), 197–215. https://doi.org/10.1207/s15327965pli1503_02
- Meng, R., Luo, Y., Liu, B., Hu, Y., & Yu, C. (2015) The Nurses' Well-Being Index and Factors Influencing This Index among Nurses in Central China: A Cross-Sectional Study. *PLoS ONE* 10(12), e0144414.
- Molero Jurado, M. D. M., Pérez-Fuentes, M. D. C., Oropesa Ruiz, N. F., Simón Márquez, M. D. M., & Gázquez Linares, J. J. (2019). Self-efficacy and emotional intelligence as predictors of perceived stress in nursing professionals. *Medicina*, 55(6), 237.
- Moore, R. C., Eyler, L. T., Mausbach, B. T., Zlatar, Z. Z., Thompson, W. K., Peavy, G., Fazeli, P. L., & Jeste, D. V. (2015). Complex interplay between health and successful aging: Role of perceived stress, resilience, and social support. *The American Journal of Geriatric Psychiatry*, 23(6), 622–632. <https://doi.org/10.1016/j.jagp.2014.08.004>

Salovey, P., & Mayer, J. D. (1990). Emotional Intelligence. *Imagination, Cognition and Personality*, 9(3), 185–211. <https://doi.org/10.2190/DUGG-P24E-52WK-6CDG>

Zhang, Y., Gan, Y., Cham, H., & Li, W. (2020). Influence of subjective well-being on work engagement among Chinese nurses: The mediating role of psychological capital. *Frontiers in Psychology*, 11, 1540.